

# **Neuroprotective Glycosides from the Root of *Iodes cirrhosa***

Maoluo Gan, Yanling Zhang, Sheng Lin, Mingtao Liu, Weixia Song, Jiachen Zi, Yongchun Yang, Xiaona Fan, Jiangong Shi,\* Jinfeng Hu, Jiandong Sun, and Naihong Chen\*

*Institute of Materia Medica, Chinese Academy of Medical Sciences and Peking Union Medical College (Key Laboratory of Bioactive Substances and Resources Utilization of Chinese Herbal Medicine, Ministry of Education), Beijing 100050, China*

shijg@imm.ac.cn

## **Supporting Information Available**


---

\* To whom correspondence should be addressed. Tel: 86-10-83154789. Fax: 86-10-63017757.  
E-mail: shijg@imm.ac.cn


## The List of Content

No	Content	Page
1	Structures of Compounds <b>1–17</b> and <b>1a–13a</b>	S4
2	<b>Figure S1.</b> Key NOE effects and HMBC correlations of compounds <b>1</b> and <b>9</b>	S5
3	<b>Figure S2.</b> CD spectra of Compounds <b>1–7</b>	S5
4	<b>Figure S3.</b> CD and UV spectra of Compounds <b>1a–7a</b>	S6
5	The $^1\text{H}$ NMR Spectrum of Compound <b>1</b> in DMSO- $d_6$	S7
6	The $^{13}\text{C}$ NMR Spectrum of Compound <b>1</b> in DMSO- $d_6$	S8
7	The NOE Difference Spectrum of Compound <b>1</b> in DMSO- $d_6$	S9
8	The $^1\text{H}$ NMR Spectrum of Compound <b>1a</b> in $\text{CDCl}_3$	S10
9	The $^1\text{H}$ NMR Spectrum of Compound <b>1b</b> in $\text{CDCl}_3$	S11
10	The $^1\text{H}$ NMR Spectrum of Compound <b>2</b> in DMSO- $d_6$	S12
11	The $^{13}\text{C}$ NMR Spectrum of Compound <b>2</b> in DMSO- $d_6$	S13
12	The $^1\text{H}$ NMR Spectrum of Compound <b>2a</b> in $\text{CDCl}_3$	S14
13	The $^1\text{H}$ NMR Spectrum of Compound <b>2b</b> in $\text{CDCl}_3$	S15
14	The $^1\text{H}$ NMR Spectrum of Compound <b>3</b> in DMSO- $d_6$	S16
15	The $^{13}\text{C}$ NMR Spectrum of Compound <b>3</b> in DMSO- $d_6$	S17
16	The $^1\text{H}$ NMR Spectrum of Compound <b>3a</b> in $\text{CDCl}_3$	S18
17	The $^1\text{H}$ NMR Spectrum of Compound <b>4</b> in DMSO- $d_6$	S19
18	The $^{13}\text{C}$ NMR Spectrum of Compound <b>4</b> in DMSO- $d_6$	S20
19	The $^1\text{H}$ NMR Spectrum of Compound <b>5</b> in DMSO- $d_6$	S21
20	The $^{13}\text{C}$ NMR Spectrum of Compound <b>5</b> in DMSO- $d_6$	S22
21	The $^1\text{H}$ NMR Spectrum of Compound <b>5a</b> $\text{CDCl}_3$	S23
22	The $^1\text{H}$ NMR Spectrum of Compound <b>6</b> in DMSO- $d_6$	S24
23	The $^{13}\text{C}$ NMR Spectrum of Compound <b>6</b> in DMSO- $d_6$	S25
24	The $^1\text{H}$ NMR Spectrum of Compound <b>6a</b> in $\text{CDCl}_3$	S26
25	The $^1\text{H}$ NMR Spectrum of Compound <b>7</b> in DMSO- $d_6$	S27
26	The $^{13}\text{C}$ NMR Spectrum of Compound <b>7</b> in DMSO- $d_6$	S28
27	The $^1\text{H}$ NMR Spectrum of Compound <b>7a</b> in $\text{CDCl}_3$	S29


28	The $^1\text{H}$ NMR Spectrum of Compound <b>7b</b> in $\text{CDCl}_3$	S30
29	The $^1\text{H}$ NMR Spectrum of Compound <b>8</b> in $\text{MeOH}-d_4$	S31
30	The $^{13}\text{C}$ NMR Spectrum of Compound <b>8</b> in $\text{MeOH}-d_4$	S32
31	The $^1\text{H}$ NMR Spectrum of Compound <b>9</b> in $\text{DMSO}-d_6$	S33
32	The $^{13}\text{C}$ NMR Spectrum of Compound <b>9</b> in $\text{DMSO}-d_6$	S34
33	The $^1\text{H}$ NMR Spectrum of Compound <b>10</b> in $\text{DMSO}-d_6$	S35
34	The $^{13}\text{C}$ NMR Spectrum of Compound <b>10</b> in $\text{DMSO}-d_6$	S36
35	The $^1\text{H}$ NMR Spectrum of Compound <b>11</b> in $\text{DMSO}-d_6$	S37
36	The $^{13}\text{C}$ NMR Spectrum of Compound <b>11</b> in $\text{DMSO}-d_6$	S38
37	The $^1\text{H}$ NMR Spectrum of Compound <b>12</b> in $\text{DMSO}-d_6$	S39
38	The $^{13}\text{C}$ NMR Spectrum of Compound <b>12</b> in $\text{DMSO}-d_6$	S40
39	The $^1\text{H}$ NMR Spectrum of Compound <b>13</b> in $\text{DMSO}-d_6$	S41
40	The $^{13}\text{C}$ NMR Spectrum of Compound <b>13</b> in $\text{DMSO}-d_6$	S42
41	<b>Table S1.</b> $^1\text{H}$ NMR data ( $\delta$ ) for compounds <b>1a</b> , <b>1b</b> , <b>2a(4a)</b> , <b>2b</b> in Different Solvents	S43
42	<b>Table S2.</b> $^1\text{H}$ NMR Data ( $\delta$ ) for Compounds <b>3a</b> , <b>5a</b> , and <b>6a</b> in Different Solvents	S44
43	<b>Table S3.</b> $^1\text{H}$ NMR Data ( $\delta$ ) for Compounds <b>7a</b> and <b>7b</b> in Different Solvents	S45
44	<b>Table S4.</b> $^1\text{H}$ NMR Data ( $\delta$ ) for Compounds <b>5</b> and <b>6</b> in Different Solvents	S46
45	<b>Table S5.</b> $^1\text{H}$ NMR Data ( $\delta$ ) for Compounds <b>7</b> in Different Solvents	S47
46	<b>Table S6.</b> $^{13}\text{C}$ NMR Data ( $\delta$ ) for Compounds <b>1a</b> , <b>2a(4a)</b> , <b>5a–7a</b> in Different Solvents	S48
47	<b>Table S7.</b> $^{13}\text{C}$ NMR Data ( $\delta$ ) for Compounds <b>2</b> and <b>5–7</b> in Different Solvents	S49
48	<b>Table S8.</b> $^1\text{H}$ NMR Data ( $\delta$ ) for Compounds <b>10</b> and <b>11</b> in Different Solvents	S50
49	<b>Table S9.</b> $^1\text{H}$ NMR Data ( $\delta$ ) for Compounds <b>12</b> and <b>13</b> in Different Solvents	S51
50	<b>Table S10.</b> $^{13}\text{C}$ NMR Data ( $\delta$ ) for Compounds <b>10–13</b> in Different Solvents	S52
51	<b>Table S11.</b> $^1\text{H}$ NMR Data ( $\delta$ ) for Compounds <b>10a</b> and <b>11a</b> in Different Solvents	S52
52	<b>Table S12.</b> $^1\text{H}$ NMR Data ( $\delta$ ) for Compounds <b>12a</b> and <b>13a</b> in Different Solvents	S53
53	<b>Table S13.</b> $^{13}\text{C}$ NMR data ( $\delta$ ) for compounds <b>10a–13a</b> in Different Solvents	S54
54	<b>Table S14.</b> $^1\text{H}$ NMR data ( $\delta$ ) for compounds <b>14–16</b> in Different Solvents	S55
55	<b>Table S15.</b> $^{13}\text{C}$ NMR data ( $\delta$ ) for Compounds <b>14–16</b> in Different Solvents	S56





**Figure S1.** Key NOE effects and HMBC correlations of compounds **1** and **9**




**Figure S2.** CD spectra of Compounds **1–7**




**Figure S3.** CD and UV spectra of Compounds **1a–7a**




The  $^1\text{H}$  NMR Spectrum of  $(-)$ -( $7S,8R,7'E$ )- 4,7,9,3',9'-Pentahydroxy-3-methoxy-8-4'-oxyneolign-7'-ene 3'- $O$ - $\beta$ -D-glucopyranoside (**1**) in  $\text{DMSO}-d_6$




The  $^{13}\text{C}$  NMR Spectrum of ( $-$ )-(7*S*,8*R*,7'*E*)- 4,7,9,3',9'-Pentahydroxy-3-methoxy-8-4'-oxyneolign-7'-ene 3'-*O*- $\beta$ -D-glucopyranoside (1) in  $\text{DMSO}-d_6$




The NOE Difference Spectrum of (*-*)-(7*S*,8*R*,7'*E*)-4,7,9,3',9'-Pentahydroxy-3-methoxy-8-4'-oxyneolign-7'-ene 3'-*O*- $\beta$ -D-glucopyranoside (1) in DMSO-*d*<sub>6</sub>

SYS-600 1H-NMR GML-C2-8b IN CDCl<sub>3</sub> 07.10.22The <sup>1</sup>H NMR Spectrum of Compound 1a in CDCl<sub>3</sub>


SYS-600 1H-NMR GML-C2-8d.in CDCL3 07.11.27




### The $^1\text{H}$ NMR Spectrum of Compound 1b in $\text{CDCl}_3$




The  $^1\text{H}$  NMR Spectrum of  $(-)(7R,8S)$ -4,7,9,3',9'-Pentahydroxy-3-methoxy-8-4'-oxyneolignan 3'- $O$ - $\beta$ -D-glucopyranoside (2) in  $\text{DMSO}-d_6$




The  $^{13}\text{C}$  NMR Spectrum of  $(-)$ -(7*R*,8*S*)- 4,7,9,3',9'-Pentahydroxy-3-methoxy-8-4'-oxyneolignan 3'-*O*- $\beta$ -D-glucopyranoside (**2**) in  $\text{DMSO}-d_6$




The <sup>1</sup>H NMR Spectrum of Compound 2a in CDCl<sub>3</sub>


SYS-600 1H-NMR GML-C2-15d in CDCl<sub>3</sub> 07.11.13The <sup>1</sup>H NMR Spectrum of Compound 2b in CDCl<sub>3</sub>




The  $^1\text{H}$  NMR Spectrum of  $(-)$ -( $7S,8R,7'E$ )-4,7,9,3',9'-Pentahydroxy-3,5'-dimethoxy-8-4'-oxyneolign-7'-ene 3'-O- $\beta$ -D-glucopyranoside (**3**) in  $\text{DMSO}-d_6$




The  $^{13}\text{C}$  Spectrum of  $(-)$ -(*7S,8R,7'E*)- 4,7,9,3',9'-Pentahydroxy-3,5'-dimethoxy-8-4'-oxyneolign-7'-ene 3'-*O*- $\beta$ -D-glucopyranoside (**3**) in DMSO- $d_6$

SYS-600 1H-NMR GML-C2-38b in CDCl<sub>3</sub> 07.11.27The <sup>1</sup>H NMR Spectrum of Compound 3a in CDCl<sub>3</sub>




The  $^1\text{H}$  NMR Spectrum of  $(-)(7R,8S)$ - 4,7,9,3',9'-Pentahydroxy-3-methoxy-8-4'-oxyneolignan 9'- $O$ - $\beta$ -D-glucopyranoside (**4**) in DMSO- $d_6$


INOVA-500 13C-NMR GML-C2-35 IN DMSO 2006.12.08




The <sup>13</sup>C NMR Spectrum of (−)-(7*R*,8*S*)-4,7,9,3',9'-Pentahydroxy-3-methoxy-8-4'-oxyneolignan 9'-*O*-β-D-glucopyranoside (4) in DMSO-*d*<sub>6</sub>




The  $^1\text{H}$  NMR Spectrum of  $(-)$ -( $7R,8S,7'E$ )-4,7,9,9'-Tetrahydroxy-3,3'-dimethoxy-8-4'-oxyneolign-7'-ene 9'- $O$ - $\beta$ -D-glucopyranoside (**5**) in  $\text{DMSO}-d_6$




The  $^{13}\text{C}$  NMR Spectrum of  $(-)(7R,8S,7'E)$ -4,7,9,9'-Tetrahydroxy-3,3'-dimethoxy-8-4'-oxyneolign-7'-ene 9'- $O$ - $\beta$ -D-glucopyranoside (**5**) in DMSO- $d_6$

SYS-600  $^1\text{H}$ -NMR GML-C2-29b IN CDCl<sub>3</sub> 07.10.22The  $^1\text{H}$  NMR Spectrum of Compound 5a CDCl<sub>3</sub>





The  $^1\text{H}$  NMR Spectrum of  $(-)$ -( $7S,8S,7'E$ )-4,7,9,9'-Tetrahydroxy-3,3'-dimethoxy-8-4'-oxyneolign-7'-ene 9'- $O$ - $\beta$ -D-glucopyranoside (**6**) in  $\text{DMSO}-d_6$

INOVA-500 13C-NMR GML-C2-33 IN DMSO 2006.12.12




The  $^{13}\text{C}$  NMR Spectrum of  $(-)$ -(7*S*,8*S*,7'*E*)-4,7,9,9'-Tetrahydroxy-3,3'-dimethoxy-8-4'-oxyneolign-7'-ene 9'-*O*- $\beta$ -D-glucopyranoside (**6**) in  $\text{DMSO}-d_6$

SYS-600 1H-NMR GML-C2-33b IN CDCl<sub>3</sub> 07.10.22The <sup>1</sup>H NMR Spectrum of Compound 6a in CDCl<sub>3</sub>




The  $^1\text{H}$  NMR Spectrum of (+)-(7*S*,8*S*)-4,7,9,9'-Tetrahydroxy-3,3'-dimethoxy-8-4'-oxyneolign 9'-*O*- $\beta$ -D-glucopyranoside (7) in DMSO- $d_6$





The  $^{13}\text{C}$  NMR Spectrum of (+)-(7*S*,8*S*)-4,7,9,9'-Tetrahydroxy-3,3'-dimethoxy-8-4'-oxyneolign 9'-*O*- $\beta$ -D-glucopyranoside (7) in DMSO- $d_6$

SYS-600 1H-NMR GML-C2-37b IN CDCl<sub>3</sub> 07.10.24




The <sup>1</sup>H NMR Spectrum of Compound 7a in CDCl<sub>3</sub>

SYS-600 1H-NMR GML-C2-37d IN CDCl<sub>3</sub> 07.10.30The <sup>1</sup>H NMR Spectrum of Compound 7b in CDCl<sub>3</sub>




The  $^1\text{H}$  NMR Spectrum of  $(-)$ -(2*R*)-1-*O*- $\beta$ -D-Glucopyranosyl-2-{2-methoxy-4-[(*E*)-formylvinyl]phenoxy}propane-3-ol (**8**) in  $\text{MeOH-}d_4$


INOVA-500 13C-NMR GML-C2-12 IN CD3OD 2006.07.18



The <sup>13</sup>C NMR Spectrum of (−)-(2*R*)-1-*O*-β-D-Glucopyranosyl-2-{2-methoxy-4-[(*E*)-formylvinyl]phenoxy}propane-3-ol (**8**) in MeOH-*d*<sub>4</sub>




The  $^1\text{H}$  NMR Spectrum of (*-*)-*Erythro*-2-(4-hydroxy-3-methoxyphenyl)-1-[2-(4-*O*- $\beta$ -D-glucopyranosyl-3-methoxyphenyl)-3-hydroxymethyl-7-methoxy-2,3-dihydrobenzofuran-5-yl]-propan-1,3-diol (9) in  $\text{DMSO}-d_6$




The  $^{13}\text{C}$  NMR Spectrum of (*-*)-*Erythro*-2-(4-hydroxy-3-methoxyphenyl)-1-[2-(4-*O*- $\beta$ -D-glucopyranosyl-3-methoxyphenyl)-3-hydroxymethyl-7-methoxy-2,3-dihydrobenzofuran-5-yl]-propan-1,3-diol (9) in DMSO- $d_6$

MP-400 H1 DMSO GML-C1-21 061116




The  $^1\text{H}$  NMR Spectrum of  $(-)(7R,8S)$ -Syringylglycerol 8- $O$ - $\beta$ -D-glucopyranoside (10) in  $\text{DMSO}-d_6$


INOVA-500  $^{13}\text{C}$ -NMR GML-C1-21 IN DMSO 2006.11.21

The  $^{13}\text{C}$  NMR Spectrum of  $(-)$ -(7*R*,8*S*)-Syringylglycerol 8-*O*- $\beta$ -D-glucopyranoside (10) in DMSO-*d*<sub>6</sub>

MP-400 H1 DMSO+D2O GML-C1-16 070929




The  $^1\text{H}$  NMR Spectrum of (+)-(7S,8S)-Syringylglycerol 8-O- $\beta$ -D-glucopyranoside (11) in DMSO- $d_6$

INOVA-500  $^{13}\text{C}$ -NMR GML-C1-16 IN DMSO


The  $^{13}\text{C}$  NMR Spectrum of (+)-(7S,8S)-Syringylglycerol 8-O- $\beta$ -D-glucopyranoside (11) in DMSO- $d_6$

MP-400 H1 DMSO GML-C1-18 061116




The  $^1\text{H}$  NMR Spectrum of  $(-)(7R,8S)$ -Guaiacylglycerol 8- $O$ - $\beta$ -D-glucopyranoside (12) in DMSO- $d_6$

INOVA-500 13C-NMR GML-C1-18 IN DMSO 2006.11.21




The  $^{13}\text{C}$  NMR Spectrum of  $(-)$ -(7*R*,8*S*)-Guaiacylglycerol 8-*O*- $\beta$ -D-glucopyranoside (12) in  $\text{DMSO}-d_6$

MP-400 H1 DMSO GML-C1-1 061206



The <sup>1</sup>H NMR Spectrum of (+)-(7*S*,8*S*)-Guaiacylglycerol 8-*O*- $\beta$ -D-glucopyranoside (13) in DMSO-*d*<sub>6</sub>

INOVA-500  $^{13}\text{C}$ -NMR GML-C1-1 IN DMSO 2006.12.08

The  $^{13}\text{C}$  NMR Spectrum of (+)-(7*S*,8*S*)-Guaiacylglycerol 8-*O*- $\beta$ -D-glucopyranoside (13) in DMSO-*d*<sub>6</sub>

**Table S1.**  $^1\text{H}$  NMR Data ( $\delta$ ) for Compounds **1a**, **1b**, **2a(4a)**, and **2b** in Different Solvents<sup>a</sup>

no.	<b>1a(Erythro)</b>			<b>1b(Erythro)</b>	<b>2a(4a) (Erythro)</b>			<b>2b(Erythro)</b>
	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub> +D <sub>2</sub> O	CD <sub>3</sub> OD	CDCl <sub>3</sub>	CDCl <sub>3</sub>	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub> +D <sub>2</sub> O	CD <sub>3</sub> OD	CDCl <sub>3</sub>	CDCl <sub>3</sub>
2	7.04 brs	6.95 brs	6.92 brs	6.94 d (1.8)	7.04 d (1.2)	6.94 d (1.8)	6.93 brs	6.94 brs
5	6.74 d (7.8)	6.69 d (7.8)	6.89 d (8.4)	6.89 d (7.8)	6.75 d (7.8)	6.69 d (8.4)	6.90 d (8.4)	6.89 d (7.8)
6	6.85 brd (7.8)	6.79 brd (7.8)	6.85 brd (8.4)	6.97 dd (7.8,1.8)	6.84 dd (7.8,1.2)	6.77 dd (8.4,1.8)	6.84 brd (8.4)	6.97 brd (7.8)
7	<b>4.91 d (4.8)</b>	<b>4.80 (overlapped)</b>	<b>5.08 d (4.2)</b>	<b>4.82 d (9.0)</b>	<b>4.91 d (4.8)</b>	<b>4.80 (overlapped)</b>	<b>5.08 d (3.6)</b>	<b>4.80 d (9.0)</b>
8	4.16 ddd (6.6,4.8,3.6)	4.15 m	4.10 ddd (5.4,4.2,3.6)	3.95 ddd (9.6,9.0,3.6)	4.06 ddd (6.6,4.8,4.2)	4.07 m	4.06 ddd (4.8,3.6,3.0)	4.12 ddd (9.0,8.4,4.8)
9a	3.83 dd (11.4,6.6)	3.80 dd (12.0,6.6)	3.96 dd (12.0,5.4)	4.19 dd (10.8,3.6)	3.81 dd (11.4,6.6)	3.78 dd (11.4,6.6)	3.94 dd (11.4,4.8)	4.17 dd (11.4,4.8)
9b	3.71 dd,(11.4,3.6)	3.70 dd (12.0,3.6)	3.80 dd (12.0,3.6)	4.16 dd (10.8,9.6)	3.68 dd (11.4,4.2)	3.67 dd (11.4,3.6)	3.77 dd (11.4,3.0)	3.95 dd (11.4,8.4)
2'	6.87 d (2.4)	6.82 brs	7.01 brs	6.89 d (1.8)	6.66 d (1.8),	6.60 d (1.8)	6.80 brs	6.66 brs
5'	6.78 d (8.4)	6.67 d (8.4)	6.91 d (8.4)	6.55 d (7.8)	6.72 d (8.4)	6.65 d (8.4)	6.86 d (7.8)	6.51 d (8.4)
6'	6.70 dd (8.4,2.4)	6.66 brd (8.4)	6.83 brd (8.4)	6.71 dd (7.8,1.8)	6.49 dd (8.4,1.8)	6.47 dd (8.4,1.8)	6.63 brd (7.8)	6.52 brd (8.4)
7'	6.42 d (15.6)	6.39 d (16.2)	6.52 d (15.6)	6.43 d (15.4)	2.51 t (7.2)	2.48 t (7.2)	2.63 t (7.2)	2.55 t (7.2)
8'	6.18 dt (15.6,5.4)	6.12 dt (16.2,6.0)	6.27 dt (15.6,5.4)	6.10 dt (15.4,6.0)	1.73 tt (7.2,6.6)	1.71 tt (7.2,6.6)	1.86 tt (7.2,6.0)	1.79 tt (7.2,6.0)
9'	4.15 d (5.4)	4.12 d (6.0)	4.30 d (5.4)	4.03 d (6.0)	3.50 t (6.6)	3.48 t (6.6)	3.67 t (6.0)	3.60 t (6.0)
3-OMe	3.76 s	3.75 s	3.88 s	3.86 s	3.77 s	3.75 s	3.89 s	3.85 s
CH <sub>3</sub> ax				1.51 s				1.51 s
CH <sub>3</sub> eq				1.64 s				1.64 s

<sup>a</sup>  $^1\text{H}$  NMR data were measured at 600 MHz. Proton coupling constants (*J*) in Hz are given in parentheses.

**Table S2.**  $^1\text{H}$  NMR Data ( $\delta$ ) for Compounds **3a**, **5a** and **6a** in Different Solvents<sup>a</sup>

no.	<b>3a (Erythro)</b>			<b>5a(Erythro)</b>			<b>6a(Threo)</b>		
	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub> +D <sub>2</sub> O	CD <sub>3</sub> OD	CDCl <sub>3</sub>	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub> +D <sub>2</sub> O	CD <sub>3</sub> OD	CDCl <sub>3</sub>	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub> +D <sub>2</sub> O	CD <sub>3</sub> OD	CDCl <sub>3</sub>
2	7.05 brs	6.96 d (1.8)	6.95 d (1.8)	7.06 d (1.8)	6.96 d (1.8)	6.96 brs	7.07 d (1.8)	6.97 d (1.8)	6.97 brs
5	6.76 d (8.4)	6.70 d (8.4)	6.89 d (8.4)	6.72 d (7.8)	6.67 d (7.8)	6.90 d (7.8)	6.74 d (7.8)	6.69 d (8.4)	6.89 d (7.8)
6	6.83 brd (8.4)	6.75 dd (8.4,1.8)	6.81 dd (8.4,1.8)	6.85 dd (7.8,1.8)	6.77 dd (7.8,1.8)	6.93 brd (7.8)	6.89 dd (7.8,1.8)	6.80 dd (8.4,1.8)	6.91 brd (7.8)
7	<b>5.06 d (3.0)</b>	<b>4.96 d (4.2)</b>	<b>5.05 d (4.2)</b>	<b>4.86 d (5.4)</b>	<b>4.76 d (6.0)</b>	<b>4.96 d (4.2)</b>	<b>4.86 d (6.6)</b>	<b>4.82 d (5.4)</b>	<b>4.95 d (8.4)</b>
8	3.99 ddd (4.2,3.0,3.0)	4.02 ddd (6.6,4.2,4.2)	4.01 ddd (6.0,4.2,3.0)	4.33 ddd, (5.4,5.4,3.6)	4.30 ddd (6.0,6.0,3.6)	4.16 ddd (5.4,4.2,3.0)	4.24 ddd (6.6,6.0,4.2)	4.23 ddd (5.4,5.4,4.2)	4.02 m
9a	3.82 dd (11.4,4.2)	3.81 dd (12.0,6.6)	3.87 dd (12.0,6.0)	3.80 dd (12.0,5.4)	3.79 dd (12.0,6.0)	3.92 dd (12.0,5.4)	3.68 dd (12.0,4.2)	3.67 dd (12.0,4.2)	3.62 brd (12.0)
9b	3.48 dd (11.4,3.0)	3.47 dd (12.0,4.2)	3.62 dd (12.0,3.0)	3.69 dd (12.0,3.6)	3.71 dd (12.0,3.6)	3.68 dd (12.0,3.0)	3.47 dd (12.0,6.0)	3.41 dd (12.0,5.4)	3.49 brd (12.0)
2'	6.61 brs	6.54 brs	6.67 d (1.8)	7.01 d (1.8)	6.94 brs	6.96 brs	7.08 d (1.8)	7.01 d (1.8)	6.97 brs
5'			6.89 d (7.8)	6.81 brs	6.88 d (8.4)	7.07 d (8.4)	6.94 d (8.4)	7.06 d (8.4)	
6'	6.57 brs	6.54 brs	6.52 d (1.8)	6.83 dd (7.8,1.8)	6.81 brs	6.83 brd (8.4)	6.86 dd (8.4,1.8)	6.85 dd (8.4,1.8)	6.94 brd (8.4)
7'	6.46 d (16.2)	6.43 brd (15.6)	6.49 brd (15.6)	6.46 brd (15.6)	6.45 brd (16.2)	6.56 d (16.2)	6.50 d (15.6)	6.48 brd (16.2)	6.56 d (16.2)
8'	6.28 dt (16.2,5.4)	6.20 dt (15.6,6.0)	6.28 dt (15.6,6.0)	6.23 dt,(15.6,5.4)	6.18 dt (16.2,5.4)	6.28 dt (16.2,5.4)	6.27 dt (15.6,5.4)	6.20 dt (16.2,6.0)	6.28 dt (16.2,6.0)
9'	4.18 d (5.4)	4.15 dd (6.0,1.2)	4.32 dd (6.0,1.2)	4.14 dd (5.4,1.2)	4.13 dd (5.4,1.2)	4.32 d (5.4)	4.17 d (5.4)	4.14 dd (6.0,1.2)	4.32 d (6.0)
3'-OMe	3.79 s	3.76 s	3.88 s	3.77 s	3.74 s	3.89 s	3.79 s	3.76 s	3.89 s
3'-OMe	3.83 s	3.78 s	3.89 s	3.81 s	3.75 s	3.90 s	3.88 s	3.82 s	3.92 s

<sup>a</sup>  $^1\text{H}$  NMR data were measured at 600 MHz. Proton coupling constants (*J*) in Hz are given in parentheses.

**Table S3.**  $^1\text{H}$  NMR Data ( $\delta$ ) for Compounds **7a** and **7b** in Different Solvents <sup>a</sup>

no.	<b>7a</b> ( <i>Threo</i> )			<b>7b</b>
	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub> +D <sub>2</sub> O	CD <sub>3</sub> OD	CDCl <sub>3</sub>	CDCl <sub>3</sub>
2	7.04 d (1.8)	6.96 d (1.8)	6.97 brs	7.20 brs
5	6.73 d (8.4)	6.69 d (7.8)	6.88 d (7.8)	6.84 d (7.8)
6	6.85 dd (8.4,1.8)	6.80 dd (7.8,1.8)	6.91 brd (7.8)	6.89 brd (7.8)
7	<b>4.86 d (6.0)</b>	<b>4.81 d (6.0)</b>	<b>4.95 d (8.4)</b>	<b>5.06 brs</b>
8	4.17 ddd (6.0,5.4,4.2)	4.13 ddd (6.0,5.4,4.2)	3.96 ddd (8.4,3.6,3.0)	4.06 d (1.8)
9a	3.67 dd (12.0,4.2)	3.65 dd (12.0,4.2)	3.61 dd (12.6,3.0)	4.18 dd (12.6,1.8)
9b	3.44 dd (12.0,5.4)	3.39 dd (12.0,5.4)	3.47 dd (12.6,3.6)	4.12 d (12.6)
2'	6.85 d (1.8)	6.79 d (1.8)	6.78 brs	6.63 brs
5'	7.00 d (8.4)	6.91 d (7.8)	7.02 d (8.4)	6.37 d (7.8)
6'	6.68 dd (8.4,1.8)	6.45 dd (7.8,1.8)	6.75 brd (8.4)	6.53 brd (7.8)
7'	2.58 t (7.8)	2.56 t (7.8)	2.67 t (7.8)	2.58 t (7.8)
8'	1.76 tt (7.8,6.0)	1.75 tt (7.8,6.0)	1.88 tt (7.8,6.6)	1.81 tt (7.8,6.0)
9	3.51 t (6.0)	3.50 t (6.0)	3.67 t (6.6)	3.62 t (6.0)
3-OMe	3.77 s	3.76 s	3.88 s	3.71 s
3'-OMe	3.83 s	3.79 s	3.90 s	3.89 s
CH <sub>3</sub> ax				1.58 s
CH <sub>3</sub> eq				1.61 s

<sup>a</sup>  $^1\text{H}$  NMR data were measured at 600 MHz. Proton coupling constants (*J*) in Hz are given in parentheses.

**Table S4.**  $^1\text{H}$  NMR Data ( $\delta$ ) for Compounds **5** and **6** in Different Solvents

no.	<b>5 (Erythro)</b>				<b>6 (Threo)</b>			
	D <sub>2</sub> O <sup>a</sup>	DMSO- <i>d</i> <sub>6</sub> <sup>b</sup>	C <sub>5</sub> D <sub>5</sub> N <sup>a</sup>	CD <sub>3</sub> OD <sup>a</sup>	D <sub>2</sub> O <sup>a</sup>	DMSO- <i>d</i> <sub>6</sub> <sup>b</sup>	C <sub>5</sub> D <sub>5</sub> N <sup>a</sup>	CD <sub>3</sub> OD <sup>a</sup>
2	6.95 brs	6.97 d (1.5)	7.59 d (1.8)	6.96 brs	7.02 brs	6.94 d (1.5)	7.58 brs	6.97 d (1.8)
5	6.79 d (7.8)	6.66 d (8.0)	7.24 d (8.4)	6.67 d (8.0)	6.87 d (7.8)	6.66 d (8.0)	7.25 d (7.8)	6.69 d (8.4)
6	6.87 brd (7.8)	6.75 dd (8.0,1.5)	7.38 dd (8.4,1.8)	6.77 brd (8.0)	6.94 brd (7.8)	6.74 dd (8.0,1.5)	7.40 brd (7.8)	6.80 dd (8.4,1.8)
7	<b>4.75 d (7.8)</b>	<b>4.69 d (5.5)</b>	<b>5.61 d (4.8)</b>	<b>4.77 d (6.0)</b>	<b>4.97 d (4.8)</b>	<b>4.69 d (4.5)</b>	<b>5.58 d (5.4)</b>	<b>4.83 d (5.4)</b>
8	4.67 m	4.30 m	5.05ddd (6.0,5.4,4.8)	4.31ddd (6.0,5.6,4.0)	4.64ddd (5.4,4.8,3.6)	4.26ddd (6.5,4.5,4.0)	4.95ddd (6.0,5.4,5.4)	4.25ddd (5.4,5.4,4.2)
9a	4.10 brd (12.0)	3.58 brs	4.55 dd (12.0,5.4)	3.79 dd (11.6,5.6)	3.84 dd (12.0,3.6)	3.57 dd (11.5,4.0)	4.42 dd (12.6,5.4)	3.67 dd (12.0,4.2)
9b	3.94 dd (12.0,6.0)		4.46 dd (12.0,6.0)	3.72 dd (11.6,4.0)	3.66 dd (12.0,5.4)	3.24 dd (11.5,6.5)	4.10 dd (12.6,6.0)	3.41 dd (12.0,5.4)
2'	6.90 brs	7.00 d (1.5)	7.02 d (1.8)	6.96 brs	7.12 brs	7.05 d (1.5)	7.03 brs	7.02 d (1.8)
5'	6.95 brs	6.91 d (8.5)	7.35 d (8.4)	6.82 brs	6.92 d (7.8)	6.96 d (8.5)	7.44 d (8.4)	6.94 d (8.4)
6'	6.95 brs	6.84 dd (8.5,1.5)	6.95 dd (8.4,1.8)	6.82 brs	6.88 brd (7.8)	6.87 dd (8.5,1.5)	6.95 brd (8.4)	6.86 dd (8.4,1.8)
7'	6.63 d (15.6)	6.53 d (16.0)	6.68 d (15.6)	6.53 d (16.2)	6.63 d (16.2)	6.54 d (16.5)	6.70 d (16.2)	6.55 d (16.2)
8'	6.25 dt (15.6,6.0)	6.20 dt (16.0,6.0)	6.35 dt (15.6,6.0)	6.18 dt (16.2,6.0)	6.27 dt (16.2,6.0)	6.23 dt (16.5,6.0)	6.37 ddd (16.2,6.6,5.4)	6.21 dt (16.2,6.0)
9'a	4.51 dd (12.6,6.0)	4.39 dd (13.0,6.0)	4.73 dd (13.2,6.0)	4.44 dd (12.8,6.0)	4.51 dd (12.0,6.0)	4.40 dd (12.0,6.0)	4.74 dd (12.6,5.4)	4.45 dd (12.6,6.0)
9'b	4.38 dd (12.6,6.0)	4.16 dd (13.0,6.0)	4.43 dd (13.2,6.0)	4.24 dd (12.8,6.0)	4.39 dd (12.0,6.0)	4.17 dd (12.0,6.0)	4.46 dd (12.6,6.6)	4.25 dd (12.6,6.0)
1''	4.55 d (7.8)	4.19 d (8.0)	4.98 d (7.8)	4.30 d (8.0)	4.54 d (7.8)	4.20 d (8.0)	4.98 d (7.8)	4.30 d (7.8)
2''	3.32 dd (8.4,7.8)	2.98 dd (8.5,8.0)	4.11 dd (8.4,7.8)	3.16 dd (8.4,8.0)	3.32 dd (8.4,7.8)	2.99 dd (8.5,8.0)	4.11 dd (8.4,7.8)	3.17 dd (8.4,7.8)
3''	3.41 dd (9.0,8.4)	3.13 dd (8.5,8.5)	4.27 dd (8.4,8.4)	3.23 dd (8.4,8.4)	3.40 dd (9.0,8.4)	3.14 dd (8.5,8.5)	4.28 dd (8.4,8.4)	3.23 dd (8.4,7.8)
4''	3.50 dd (9.0,9.0)	3.04 dd (9.0,8.5)	4.27 dd (8.4,8.4)	3.30 dd (8.4,8.0)	3.49 dd (9.0,9.0)	3.04 dd (9.0,8.5)	4.28 dd (8.4,8.4)	3.30 dd (8.4,7.8)
5''	3.46 m	3.06 m	3.97 m	3.23 m	3.44 m	3.09 m	3.97 m	3.22 m
6''a	3.93 brd (12.0)	3.66 brd (12.0)	4.58 brd (12.6)	3.82 brd (11.6)	3.92 brd (12.0)	3.67 dd (12.0,1.5)	4.58 brd (10.8)	3.82 dd (12.0,2.4)
6''b	3.74 dd (12.0,5.4)	3.44 dd (12.0,5.0)	4.41 dd (12.6,4.8)	3.62 dd (11.6,5.2)	3.73 dd (12.0,6.0)	3.44 dd (12.0,5.5)	4.40 dd (10.8,4.8)	3.62 dd (12.0,5.4)
3'-OMe	3.69 s	3.71 s	3.68 s	3.74 s	3.78 s	3.70 s	3.71 s	3.76 s
3'-OMe	3.69 s	3.72 s	3.72 s	3.74 s	3.88 s	3.78 s	3.72 s	3.82 s

<sup>a</sup>  $^1\text{H}$  NMR data were measured at 600 MHz. <sup>b</sup>  $^1\text{H}$  NMR data were measured at 500 MHz. Proton coupling constants (*J*) in Hz are given in parentheses.

**Table S5.**  $^1\text{H}$  NMR Data ( $\delta$ ) for Compounds **7** in Different Solvents

no.	<b>7 (Threo)</b>			
	D <sub>2</sub> O <sup>a</sup>	DMSO- <i>d</i> <sub>6</sub> <sup>b</sup>	C <sub>5</sub> D <sub>5</sub> N <sup>a</sup>	CD <sub>3</sub> OD <sup>a</sup>
2	7.00 brs	6.96 d (1.5)	7.56 brs	6.96 d (1.8)
5	6.84 d (7.8)	6.67 d (8.0)	7.24 d (8.4)	6.69 d (8.4)
6	6.89 brd (7.8)	6.75 dd (8.0,1.5)	7.39 brd (8.4)	6.80 dd (8.4,1.8)
7	<b>4.96 d (5.4)</b>	<b>4.68 d (4.5)</b>	<b>5.58 d (6.0)</b>	<b>4.82 d (6.0)</b>
8	4.59 ddd (6.0,5.4,4.2)	4.16 ddd (5.5,4.5,4.0)	4.88 ddd (6.0,6.0,5.4)	4.15 ddd (6.0,5.4,4.2)
9a	3.84 dd (12.0,4.2)	3.55 dd (11.0,4.0)	4.39 dd (12.0,5.4)	3.65 dd (12.0,4.2)
9b	3.66 dd (12.0,6.0)	3.22 dd (11.0,5.5)	4.06 dd (12.0,6.0)	3.40 dd (12.0,5.4)
2'	6.90 brs	6.81 d (1.5)	6.90 d (1.8)	6.82 d (1.8)
5'	6.82 d (8.4)	6.90 d (8.5)	7.43 d (8.4)	6.90 d (7.8)
6'	6.71 brd (8.4)	6.66 dd (8.5,1.5)	6.76 dd (8.4,1.8)	6.67 dd (7.8,1.8)
7'	2.62 t (7.2)	2.56 t (7.5)	2.68 m	2.62 m
8'	1.88 m	1.78 m	1.96 m	1.83 m
9'a	3.90 dt (12.0,6.0)	3.79 dt ((10.0,7.5)	4.13 dt (12.6,6.6)	3.86 dt (10.4,6.6)
9'b	3.61 dt (12.0,6.0)	3.39 dt (10.0,6.0)	3.66 dt (12.6,6.6)	3.47 dt (10.4,6.6)
1''	4.41 d (8.4)	4.09 d (8.0)	4.83 d (7.8)	4.19 d (7.2)
2''	3.29 dd (9.0,8.4)	2.95 dd (8.0,8.0)	4.06 dd (8.4,7.8)	3.14 dd (7.8,7.2)
3''	3.40 dd (9.0,9.0)	3.14 dd (8.0,8.0)	4.24 dd (8.4,8.4)	3.23 dd (8.4,7.8)
4''	3.50 dd (9.0,9.0)	3.02 dd (8.0,8.5)	4.24 dd (8.4,8.4)	3.30 dd (9.0,8.4)
5''	3.42 m	3.07 m	3.94 m	3.19 m
6''a	3.90 brd (12.0)	3.63 dd (12.0,2.0)	4.55 brd (11.2)	3.80 dd (12.0,2.4)
6''b	3.73 dd (12.0,5.4)	3.41 dd (12.0,6.0)	4.36 dd (11.2,4.8)	3.60 ddd (12.0,6.0)
3'-OMe	3.77 s	3.68 s	3.69 s	3.76 s
3'-OMe	3.84 s	3.70 s	3.74 s	3.82 s

<sup>a</sup>  $^1\text{H}$  NMR data were measured at 600 MHz. <sup>b</sup>  $^1\text{H}$  NMR data were measured at 500 MHz. Proton coupling constants (*J*) in Hz are given in parentheses.

**Table S6.**  $^{13}\text{C}$  NMR Data ( $\delta$ ) for Compounds **1a**, **2a(4a)**, **5a–7a** in Different Solvents<sup>a</sup>

no.	<b>1a(Erythro)</b>		<b>2a(4a,Erythro)</b>		<b>5a(Erythro)</b>			<b>6a(Threo)</b>			<b>7a(Threo)</b>		
	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub>	CD <sub>3</sub> OD	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub>	CD <sub>3</sub> OD	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub>	CD <sub>3</sub> OD	CDCl <sub>3</sub>	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub>	CD <sub>3</sub> OD	CDCl <sub>3</sub>	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub>	CD <sub>3</sub> OD	CDCl <sub>3</sub>
1	133.5	133.8	133.5	133.8	132.9	134.1	131.7	133.8	133.8	131.4	133.7	133.8	131.5
2	111.4	111.7	111.4	111.7	111.4	111.9	109.8	111.4	111.7	109.8	111.4	111.7	109.4
3	146.9	148.8	146.9	148.8	148.0	148.7	146.6	148.0	148.8	146.7	148.0	148.8	146.6
4	145.4	147.2	145.4	147.1	146.7	147.0	145.1	146.8	147.2	145.6	146.6	147.2	145.5
5	115.3	115.8	115.3	115.8	115.1	115.7	114.3	115.1	115.8	114.3	115.2	115.8	114.3
6	120.5	120.8	120.5	120.6	120.2	120.7	120.1	120.3	120.8	120.2	120.3	120.8	120.2
7	<b>73.7</b>	<b>74.0</b>	<b>73.8</b>	<b>74.0</b>	<b>73.8</b>	<b>74.1</b>	<b>72.8</b>	<b>73.8</b>	<b>74.0</b>	<b>74.0</b>	<b>73.3</b>	<b>74.2</b>	<b>74.0</b>
8	<b>88.1</b>	<b>87.3</b>	<b>88.4</b>	<b>87.8</b>	<b>86.7</b>	<b>86.2</b>	<b>87.4</b>	<b>88.5</b>	<b>87.2</b>	<b>89.5</b>	<b>87.4</b>	<b>87.8</b>	<b>89.6</b>
9	61.7	62.0	61.7	62.0	61.8	62.2	60.8	61.9	61.9	61.1	61.3	61.9	61.0
1'	134.1	133.7	139.1	138.7	132.9	133.1	133.1	133.0	133.2	133.1	137.4	138.2	138.2
2'	114.4	114.6	117.0	117.2	111.0	111.4	108.6	110.8	111.3	109.4	113.5	113.9	112.4
3'	150.2	149.6	153.6	149.5	151.9	151.9	151.6	151.7	151.8	151.2	150.9	151.7	151.1
4'	148.1	147.3	150.1	145.7	148.6	149.0	147.6	149.2	149.3	147.3	147.2	147.6	145.6
5'	118.5	119.3	119.9	120.2	119.4	118.9	119.0	119.7	118.9	120.1	118.7	119.7	121.0
6'	121.0	119.5	121.3	120.8	120.5	121.0	120.8	120.5	120.8	120.8	121.5	122.0	121.3
7'	129.7	128.4	32.3	32.5	129.6	128.5	128.1	129.7	128.6	128.2	32.3	32.7	31.8
8'	129.8	131.5	35.5	35.5	129.8	131.5	130.5	129.8	131.4	130.5	35.3	35.6	34.2
9'	63.3	63.8	61.7	62.3	63.3	63.8	63.6	63.3	63.8	63.6	61.5	62.2	62.1
3'-OMe	56.2	56.4	56.2	56.4	56.2	56.3	56.0	56.2	56.3	55.9	56.1	56.3	55.9
3'-OMe					56.3	56.5	56.0	56.3	56.6	56.0	56.2	56.5	56.0
$\Delta\delta_{\text{C}8-\text{C}7}$	<b>14.4</b>	<b>13.3</b>	<b>14.6</b>	<b>13.8</b>	<b>12.9</b>	<b>12.1</b>	<b>14.6</b>	<b>14.7</b>	<b>13.2</b>	<b>15.5</b>	<b>14.1</b>	<b>13.6</b>	<b>15.6</b>

<sup>a</sup>  $^{13}\text{C}$  NMR data ( $\delta$ ) were measured at 150 MHz. The assignments were based on DEPT,  $^1\text{H}$ - $^1\text{H}$  COSY, HMQC, and HMBC experiments.

**Table S7.**  $^{13}\text{C}$  NMR Data ( $\delta$ ) for Compounds **2** and **5–7** in Different Solvents<sup>a</sup>

no.	<b>2(Erythro)</b>		<b>5(Erythro)</b>				<b>6(Threo)</b>				<b>7(Threo)</b>			
	CD <sub>3</sub> OD	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD	
1	134.2	134.9	133.1	134.8	134.1	135.3	132.6	134.1	133.8	135.4	133.0	134.1	133.8	
2	111.7	112.6	111.4	112.2	111.9	112.9	111.0	111.9	111.7	113.7	111.0	111.9	111.7	
3	148.8	149.9	146.9	148.7	148.7	150.1	147.0	148.5	148.8	150.0	147.0	148.5	148.8	
4	147.0	147.2	145.5	147.7	147.0	147.4	145.8	147.6	147.2	147.3	145.4	147.6	147.2	
5	115.9	117.7	114.5	116.2	115.7	118.1	114.7	116.1	115.8	117.9	114.6	116.1	115.8	
6	120.9	122.6	119.5	120.4	120.8	122.6	119.0	120.6	120.7	122.5	119.0	120.7	120.8	
7	<b>73.4</b>	<b>75.4</b>	<b>71.6</b>	<b>73.7</b>	<b>74.1</b>	<b>75.4</b>	<b>70.9</b>	<b>73.3</b>	<b>74.0</b>	<b>75.3</b>	<b>71.0</b>	<b>73.5</b>	<b>74.1</b>	
8	<b>87.1</b>	<b>86.4</b>	<b>83.7</b>	<b>86.3</b>	<b>86.2</b>	<b>86.1</b>	<b>84.3</b>	<b>87.2</b>	<b>87.1</b>	<b>86.4</b>	<b>84.8</b>	<b>87.8</b>	<b>87.7</b>	
9	61.7	64.4	60.1	61.9	62.2	63.6	60.1	61.8	61.9	63.6	60.1	61.8	61.9	
1'	138.3	133.6	129.5	131.4	132.8	133.4	129.6	131.2	132.9	139.1	134.8	136.3	138.2	
2'	120.2	114.2	109.9	111.0	111.4	113.9	109.8	110.7	111.3	115.6	112.9	113.5	114.1	
3'	149.7	152.0	149.7	151.3	151.9	151.9	149.6	151.0	151.7	151.7	149.5	151.0	151.7	
4'	147.9	150.0	147.9	149.3	149.1	150.1	148.2	149.6	149.4	148.1	146.4	147.7	147.6	
5'	120.4	119.3	115.4	117.6	118.8	118.2	115.3	117.5	118.7	118.7	116.0	118.4	119.6	
6'	124.4	123.9	119.3	120.8	121.0	122.9	119.4	120.2	121.0	123.8	120.2	121.2	122.1	
7'	32.4	136.4	131.4	132.5	133.6	136.3	131.4	132.3	133.6	33.6	31.1	32.1	32.7	
8'	35.3	126.1	124.0	125.0	125.2	126.2	124.1	124.9	125.3	33.5	31.2	32.2	32.7	
9'	62.2	73.2	68.7	70.2	70.8	73.2	68.7	70.0	70.9	72.3	67.9	69.0	69.9	
1''	103.7	103.8	102.1	104.1	103.2	103.9	102.1	104.0	103.3	105.2	103.0	104.9	104.5	
2''	75.1	76.0	73.5	75.5	75.1	76.0	73.5	75.3	75.2	76.0	73.5	75.3	75.2	
3''	77.9	78.7	76.7	78.8	78.0	78.7	76.7	78.6	78.0	78.7	76.7	78.5	78.0	
4''	71.5	72.5	70.1	71.9	71.7	72.5	70.1	71.7	71.7	72.5	70.0	71.7	71.7	
5''	78.2	78.8	76.9	78.9	78.1	78.8	76.9	78.6	78.1	78.7	76.8	78.6	78.1	
6''	62.6	63.6	61.1	63.0	62.8	63.6	61.1	62.8	62.8	63.6	61.1	62.8	62.8	
3'-OMe	56.5	58.2	55.4	56.0	56.3	58.5	55.4	55.8	56.3	58.5	55.4	55.8	56.4	
3'-OMe		58.6	55.6	56.0	56.5	58.7	55.6	55.8	56.5	58.6	55.6	56.0	56.6	
$\Delta\delta_{\text{C}8-\text{C}7}$	<b>13.7</b>	<b>11.0</b>	<b>12.1</b>	<b>12.6</b>	<b>12.1</b>	<b>10.7</b>	<b>13.4</b>	<b>13.9</b>	<b>13.1</b>	<b>11.1</b>	<b>13.8</b>	<b>14.3</b>	<b>13.6</b>	

<sup>a</sup>  $^{13}\text{C}$  NMR data ( $\delta$ ) were measured at 125 MHz. The assignments were based on DEPT,  $^1\text{H}$ - $^1\text{H}$  COSY, HMQC, and HMBC experiments.

**Table S8.**  $^1\text{H}$  NMR Data ( $\delta$ ) for Compounds **10** and **11** in Different Solvents<sup>a</sup>

no.	<b>10 (Erythro)</b>				<b>11 (Threo)</b>			
	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD
2,6	6.82 s	6.64 s	7.28 s	6.68 s	6.81 s	6.63 s	7.16 s	6.65 s
7	<b>4.82 d (6.0)</b>	<b>4.63 d (4.4)</b>	<b>5.53 d (4.0)</b>	<b>4.72 d (4.4)</b>	<b>4.80 (overlap)</b>	<b>4.53 d (6.0)</b>	<b>5.35 d (7.6)</b>	<b>4.63 d (6.4)</b>
8	4.08 m	3.75 m	4.70 m	3.86 m	4.07 m	3.61 m	4.42 m	3.76 m
9a	3.78 dd (11.2,2.8)	3.42 dd (11.2,4.4)	4.35 brd (11.6)	3.58 dd (12.0,4.8)	3.73 brd (11.6)	3.34 dd (11.2,3.6)	4.13 m	3.51 dd (11.6,3.6)
9b	3.72 dd (11.2,5.6)	3.29 dd (11.2,4.8)	4.25 dd (11.6,3.6)	3.52 dd (12.0,3.2)	3.50 dd (11.6,5.6)	3.19 dd (11.2,6.0)	3.93 m	3.35 dd (11.6,5.6)
1'	4.27 d (8.0)	4.25 d (8.0)	5.23 d (7.6)	4.27 d (7.6)	4.43 d (7.2)	4.23 d (7.6)	5.28 d (7.6)	4.31 d (7.2)
2'	3.26 dd (8.0,8.0)	3.00 dd (8.8,8.0)	4.08 dd (8.0,7.6)	3.18 dd (8.4,7.6)	3.37 m	3.04 dd (8.4,7.6)	4.13 m	3.16 dd (8.0,7.2)
3'	3.37 m	3.10 dd (8.8,8.8)	4.21 m	3.25 dd (8.4,8.0)	3.37 m	3.12 dd (8.4,8.0)	4.21 m	3.25 dd (8.8,8.0)
4'	3.37 m	3.02 dd (8.8,8.8)	4.23 m	3.28 dd (8.4,8.0)	3.37 m	3.05 dd (8.4,8.0)	4.23 m	3.27 dd (8.8,8.8)
5'	3.37 m	3.10 m	3.96 m	3.20 m	3.37 m	3.08 m	3.93 m	3.18 m
6'a	3.90 brd (11.6)	3.66 dd (11.6,1.2)	4.51 brd (11.2)	3.78 brd (12.0)	3.90 brd (12.0)	3.64 brd (12.0)	4.49 brd (11.6)	3.76 brd (11.6)
6'b	3.76 dd (11.6,6.0)	3.38 dd (11.6,5.2)	4.32 dd (11.2,5.6)	3.59 dd (12.0,4.8)	3.70 brd (12.0)	3.39 dd (12.0,5.6)	4.31 dd (11.6,5.2)	3.59 dd (11.6,5.6)
OMe	3.89 s	3.73 s	3.75 s	3.80 s	3.90 s	3.73 s	3.73 s	3.80 s

<sup>a</sup>  $^1\text{H}$  NMR data were measured at 400 MHz. Proton coupling constants (*J*) in Hz are given in parentheses.

**Table S9.**  $^1\text{H}$  NMR Data ( $\delta$ ) for Compounds **12** and **13** in Different Solvents<sup>a</sup>

no.	<b>12 (Erythro)</b> <sup>b</sup>			<b>13 (Threo)</b>			
	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD
2	7.11 s	6.97 d (1.2)	7.60 brs	7.09 s	6.95 d (1.6)	7.50 s	6.95 d (1.6)
5	6.95 s	6.67 d (8.0)	7.21 d (8.0)	6.95 s	6.68 d (8.0)	7.21 d (8)	6.71 d (8.0)
6	6.95 s	6.73 dd (8.0,1.2)	7.33 brd (8.0)	6.95 s	6.73 dd (8.0,1.6)	7.27 brd (8.0)	6.76 dd (8.0,1.6)
7	<b>4.83 d (6.0)</b>	<b>4.63 d (4.0)</b>	<b>5.50 d (4.0)</b>	<b>4.83 d (6.8)</b>	<b>4.53 d (6.4)</b>	<b>5.33 d (7.6)</b>	<b>4.61 d (6.8)</b>
8	4.07 m	3.72 m	4.66 m	4.05 m	3.60 m	4.42 m	3.74 m
9a	3.78 m	3.39 dd (11.2,5.2)	4.28 dd (11.6,4.0)	3.67 dd (12.0,3.2)	3.33 brd (11.2)	4.09 brd (11.6)	3.48 dd (12.0,3.2)
9b	3.72 m	3.25 dd (11.2,4.8)	4.19 dd (11.6,5.6)	3.47 dd (12.0,5.6)	3.17 dd (11.2,5.6)	3.91 dd (11.6,6.4)	3.32 dd (12.0,6.0)
1'	4.33 d (8.0)	4.26 d (7.6)	5.22 d (7.6)	4.50 d (7.6)	4.26 d (8.0)	5.31 d (7.6)	4.34 d (7.6)
2'	3.26 dd (8.0,7.6)	3.00 dd (8.8,7.6)	4.08 dd (8.0,7.6)	3.37 dd (8.8,7.6)	3.04 dd (8.0,8.0)	4.15 dd (8.0,7.6)	3.20 dd (8.0,7.6)
3'	3.38 m	3.13 dd (8.8,8.8)	4.19 m	3.40 m	3.13 dd (8.8,8.0)	4.23 dd (8.8,8.0)	3.26 dd (8.0,8.0)
4'	3.38 m	3.02 dd (8.8,8.4)	4.21 m	3.40 m	3.06 dd (8.8,8.0)	4.25 dd (8.8,8.0)	3.28 dd (8.0,8.0)
5'	3.38 m	3.10 m	3.96 m	3.40 m	3.09 m	3.98 m	3.20 m
6'a	3.90 brd (12.0)	3.64 brd (12.0)	4.50 dd (11.6,2.0)	3.90 brd (12.0)	3.65 brd (11.6)	4.52 brd (11.2)	3.78 dd (12.0,2.0)
6'b	3.73 m	3.39(12.0,6.4)	4.32 brd (11.6,5.6)	3.73 dd (12.0,4.0)	3.40 dd (11.6,6.0)	4.34 dd (11.2,5.2)	3.59 dd (12.0,5.2)
OMe	3.91 s	3.74 s	3.71 s	3.90 s	3.74 s	3.70 s	3.81 s

<sup>a</sup>  $^1\text{H}$  NMR data were measured at 400 MHz. Proton coupling constants (*J*) in Hz are given in parentheses.

<sup>b</sup> The  $^{13}\text{C}$  NMR data of **12** in CD<sub>3</sub>OD were not measured due to a loss of the compound.

**Table S10.**  $^{13}\text{C}$  NMR Data ( $\delta$ ) for Compounds **10–13** in Different Solvents<sup>a</sup>

no.	<b>10 (Erythro)</b>				<b>11 (Threo)</b>				<b>12 (Erythro)<sup>b</sup></b>			<b>13 (Threo)</b>			
	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD
1	134.8	131.9	133.2	132.9	134.6	131.8	132.3	132.8	135.4	132.7	134.1	135.2	132.5	133.2	133.5
2	107.6	104.6	105.7	105.2	107.2	104.3	105.7	105.2	114.2	111.5	112.1	113.9	111.0	111.7	111.5
3	150.6	147.4	148.9	149.1	150.6	147.5	149.0	149.2	150.3	147.0	148.4	150.3	147.2	148.6	149.0
4	136.5	134.3	136.7	135.9	136.5	134.5	137.2	136.2	147.6	145.4	147.3	147.6	145.6	147.8	147.4
5	150.6	147.4	148.9	149.1	150.6	147.5	149.0	149.2	118.2	114.6	115.9	118.3	114.8	116.1	116.0
6	107.6	104.6	105.7	105.2	107.2	104.3	105.7	105.2	123.0	119.4	120.6	122.8	119.4	120.9	120.8
7	<b>75.6</b>	<b>72.6</b>	<b>74.4</b>	<b>74.6</b>	<b>76.3</b>	<b>72.3</b>	<b>74.9</b>	<b>75.0</b>	<b>75.3</b>	<b>72.5</b>	<b>74.2</b>	<b>76.4</b>	<b>72.3</b>	<b>74.8</b>	<b>75.0</b>
8	<b>85.7</b>	<b>84.6</b>	<b>86.6</b>	<b>85.5</b>	<b>87.0</b>	<b>86.3</b>	<b>89.4</b>	<b>87.2</b>	<b>86.0</b>	<b>84.9</b>	<b>86.7</b>	<b>87.5</b>	<b>86.7</b>	<b>89.6</b>	<b>87.7</b>
9	64.1	61.1	62.8	62.6	64.4	61.3	63.0	63.2	64.0	61.1	62.7	64.4	61.3	63.0	63.2
1'	104.9	103.1	105.0	104.0	105.7	103.8	106.2	105.2	105.0	103.3	105.1	105.9	104.0	106.2	105.3
2'	76.1	73.8	75.6	75.3	76.1	73.9	75.7	75.5	76.1	73.9	75.5	76.1	73.9	75.7	75.5
3'	78.4	76.4	78.3	77.8	78.5	76.3	78.4	77.9	78.4	76.4	78.2	78.5	76.3	78.5	77.9
4'	72.4	70.1	71.6	71.5	72.4	70.1	71.6	71.4	72.4	70.1	71.5	72.4	70.1	71.6	71.4
5'	78.8	76.9	78.6	78.1	78.8	76.8	78.6	78.1	78.7	76.9	78.6	78.7	76.8	78.6	78.0
6'	63.5	61.0	62.5	62.6	63.5	61.1	62.5	62.5	63.5	61.0	62.5	63.5	61.0	62.5	62.5
OMe	59.2	55.9	56.3	56.7	59.2	55.9	56.3	56.7	58.8	55.6	55.8	58.8	55.5	55.8	56.4
$\Delta\delta_{\text{C}8-\text{C}7}$	<b>10.1</b>	<b>12.0</b>	<b>12.2</b>	<b>10.9</b>	<b>10.7</b>	<b>14.0</b>	<b>14.5</b>	<b>12.2</b>	<b>10.7</b>	<b>12.4</b>	<b>12.5</b>	<b>11.1</b>	<b>14.4</b>	<b>14.8</b>	<b>12.7</b>

<sup>a</sup>  $^{13}\text{C}$  NMR data were measured at 125 MHz. The assignments were based on DEPT, <sup>1</sup>H-<sup>1</sup>H COSY, HMQC, and HMBC experiments.

<sup>b</sup> The  $^{13}\text{C}$  NMR data of **12** in CD<sub>3</sub>OD were not measured due to the loss of the compound.

**Table S11.**  $^1\text{H}$  NMR Data ( $\delta$ ) for Compounds **10a** and **11a** in Different Solvents<sup>a</sup>

no.	<b>10a (Erythro)</b>				<b>11a (Threo)</b>			
	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub> +D <sub>2</sub> O	CD <sub>3</sub> OD	C <sub>5</sub> D <sub>5</sub> N	DMSO- <i>d</i> <sub>6</sub>	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub> +D <sub>2</sub> O	CD <sub>3</sub> OD	C <sub>5</sub> D <sub>5</sub> N	DMSO- <i>d</i> <sub>6</sub>
2,6	6.69 s	6.64 s	7.21 s	6.57 s	6.67 s	6.63 s	7.19 s	6.55 s
7	4.55 d (5.6)	4.46 d (6.0)	5.32 d (5.2)	4.32 brs	4.53 d (6.0)	4.47 d (6.4)	5.33 d (6.4)	4.37 d (5.2)
8	3.68 m	3.67 m	4.51 m	3.48 m	3.62 m	3.61 m	4.42 m	3.44 m
9a	3.59 dd (11.2,4.0)	3.60 dd (11.2,4.0)	4.47 dd (11.2,4.0)	3.46 dd (11.2,4.0)	3.46 dd (11.2,4.4)	3.44 dd (11.2,4.0)	4.27 dd (11.2,4.0)	3.30 dd (11.2,4.8)
9b	3.54 dd (11.2,6.4)	3.53 dd (11.2,6.4)	4.45 dd (11.2,6.4)	3.36 dd (11.2,6.4)	3.35 dd (11.2,6.0)	3.31 dd (11.2,6.0)	4.12 dd (11.2,6.4)	3.13 dd (11.2,6.4)
OMe	3.80 s	3.79 s	3.71 s	3.72 s	3.78 s	3.79 s	3.72 s	3.70 s

<sup>a</sup>  $^1\text{H}$  NMR data were measured at 400 MHz. Proton coupling constants (*J*) in Hz are given in parentheses.

**Table S12.**  $^1\text{H}$  NMR Data ( $\delta$ ) for Compounds **12a** and **13a** in Different Solvents.<sup>a</sup>

no.	<b>12a (Erythro)</b>				<b>13a (Threo)</b>			
	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub> +D <sub>2</sub> O	CD <sub>3</sub> OD	C <sub>5</sub> D <sub>5</sub> N	DMSO- <i>d</i> <sub>6</sub>	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub> +D <sub>2</sub> O	CD <sub>3</sub> OD	C <sub>5</sub> D <sub>5</sub> N	DMSO- <i>d</i> <sub>6</sub>
2	6.98 brs	6.95 d (1.6)	7.50 brs	6.87 brs	6.99 brs	6.93 brs	7.48 brs	6.88 brs
5	6.72 d (8.0)	6.70 d (8.0)	7.24 d (8.0)	6.65 d (8.0)	6.74 d (8.0)	6.69 d (8.0)	7.24 d (8.0)	6.66 d (8.0)
6	6.79 brd (8.0)	6.76 dd (8.0,1.6)	7.37 brd (8.0)	6.68 brd (8.0)	6.79 brd (8.0)	6.74 brd (8.0)	7.33 brd (8.0)	6.68 brd (8.0)
7	4.54 d (6.0)	4.47 d (6.4)	5.30 d (6.0)	4.31 d (6.0)	4.53 d (6.0)	4.46 d (6.4)	5.30 d (6.0)	4.36 d (5.2)
8	3.68 m	3.68 m	4.50 m	3.48 m	3.61 m	3.61 m	4.39 m	3.42 m
9a	3.60 dd (11.2,4.0)	3.60 dd (11.2,4.0)	4.45 dd (11.2,4.0)	3.43 dd (11.2,4.4)	3.46 dd (11.2,4.0)	3.42 dd (11.2,4.0)	4.22 dd (11.2,4.0)	3.29 dd (10.8,4.4)
9b	3.56 dd (11.2,6.4)	3.52 dd (11.2,6.4)	4.40 dd (11.2,6.4)	3.33 dd (11.2,5.6)	3.34 dd (11.2,6.0)	3.29 dd (11.2,6.4)	4.09 dd (11.2,6.4)	3.11 dd (10.8,6.4)
OMe	3.79 s	3.80	3.65 s	3.72 s	3.80 s	3.80 s	3.66 s	3.72 s

<sup>a</sup>  $^1\text{H}$  NMR data were measured at 400 MHz. Proton coupling constants (*J*) in Hz are given in parentheses.

**Table S13.**  $^{13}\text{C}$  NMR data ( $\delta$ ) for compounds **10a–13a** in Different Solvents<sup>a</sup>

no.	<b>10a (Erythro)</b>				<b>11a (Threo)</b>				<b>12a (Erythro)</b>				<b>13a (Threo)</b>			
	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub>	CD <sub>3</sub> OD	C <sub>5</sub> D <sub>5</sub> N	DMSO- <i>d</i> <sub>6</sub>	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub>	CD <sub>3</sub> OD	C <sub>5</sub> D <sub>5</sub> N	DMSO- <i>d</i> <sub>6</sub>	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub>	CD <sub>3</sub> OD	C <sub>5</sub> D <sub>5</sub> N	DMSO- <i>d</i> <sub>6</sub>	Me <sub>2</sub> CO- <i>d</i> <sub>6</sub>	CD <sub>3</sub> OD	C <sub>5</sub> D <sub>5</sub> N	DMSO- <i>d</i> <sub>6</sub>
1	134.0	134.1	134.7	133.5	133.9	134.1	134.5	133.4	135.1	134.8	135.3	134.4	134.9	134.8	135.1	134.3
2	105.3	105.5	105.8	104.5	105.1	105.2	105.5	104.0	111.2	111.9	111.9	111.3	111.1	111.5	111.7	110.9
3	148.3	149.0	148.8	147.3	148.4	149.1	149.0	147.4	147.9	148.8	148.4	146.9	148.0	148.9	148.5	146.9
4	136.0	136.0	136.7	134.1	136.0	136.0	136.2	134.1	146.5	147.0	147.3	145.2	146.7	147.1	147.4	145.2
5	148.3	149.0	148.8	147.3	148.4	149.1	149.0	147.4	115.1	115.7	116.0	114.5	115.2	115.9	116.1	114.6
6	105.3	105.5	105.8	104.5	105.1	105.2	105.5	104.0	120.5	121.0	120.9	119.6	120.3	120.7	120.5	119.0
7	<b>76.2</b>	<b>76.3</b>	<b>76.5</b>	<b>74.0</b>	<b>75.0</b>	<b>75.6</b>	<b>75.1</b>	<b>72.9</b>	<b>76.106</b>	<b>76.1</b>	<b>76.2</b>	<b>73.9</b>	<b>74.8</b>	<b>75.4</b>	<b>74.9</b>	<b>72.7</b>
8	<b>76.4</b>	<b>76.7</b>	<b>77.0</b>	<b>75.3</b>	<b>77.3</b>	<b>77.6</b>	<b>77.8</b>	<b>75.8</b>	<b>76.106</b>	<b>76.7</b>	<b>77.0</b>	<b>75.3</b>	<b>77.3</b>	<b>77.6</b>	<b>77.8</b>	<b>75.8</b>
9	64.4	64.5	65.0	62.9	64.0	64.3	64.4	62.5	64.4	64.5	65.0	63.1	63.9	64.3	64.3	62.4
OMe	56.5	56.7	56.2	55.9	56.6	56.8	56.3	55.8	56.1	56.3	55.7	55.5	56.2	56.3	55.8	55.5
$\Delta\delta_{\text{C}8-\text{C}7}$	<b>0.2</b>	<b>0.4</b>	<b>0.5</b>	<b>1.3</b>	<b>2.3</b>	<b>2.0</b>	<b>2.7</b>	<b>2.9</b>	<b>0.000</b>	<b>0.6</b>	<b>0.8</b>	<b>1.4</b>	<b>2.5</b>	<b>2.2</b>	<b>2.9</b>	<b>3.1</b>

<sup>a</sup>  $^{13}\text{C}$  NMR data were measured at 125 MHz. The assignments were based on DEPT,  $^1\text{H}$ - $^1\text{H}$  COSY, HMQC, and HMBC experiments.

**Table S14.**  $^1\text{H}$  NMR data ( $\delta$ ) for compounds **14–16** in Different Solvents<sup>a</sup>

no.	<b>14 (Erythro)</b>				<b>15 (Threo)</b>				<b>16 (Threo)</b>			
	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD
2	7.08 s	6.89 d (1.2)	7.45 brs	6.95 brs	7.08 s	6.94 d (2.0)	7.53 brs	6.95 d (2.0)	6.75 s	6.62 s	7.28 s	6.65 s
5	6.93 s	6.67 d (8.0)	7.20 d (8.0)	6.69 d (8.0)	6.96 s	6.68 d (8.0)	7.21 d (8.0)	6.69 d (8.0)				
6	6.93 s	6.71 dd (8.0,1.2)	7.29 brd (8.0)	6.76 brd (8.0)	6.96 s	6.75 dd (8.0,2.0)	7.36 brd (8.0)	6.77 dd (8.0,2.0)	6.75 s	6.62 s	7.28 s	6.65 s
7	<b>4.65 d (7.2) 4.36 d (4.0)</b>	<b>5.26 d (6.4) 4.50 d (6.4)</b>			<b>4.68 d (6.8) 4.48 d (5.2)</b>	<b>5.35 d (5.6) 4.51 d (6.4)</b>			<b>4.65 d (6.8) 4.48 d (5.2)</b>	<b>5.42 d (5.6) 4.54 d (6.0)</b>		
8	4.07 m	3.67 m	4.58 m	3.85 m	4.02 m	3.63 m	4.44 m	3.78 m	3.98 m	3.64 m	4.45 m	3.77 m
9a	4.13 dd (12.0,1.2)	3.86 dd (10.4,2.8)	4.73 dd (10.4,2.4)	3.99 dd (10.4,2.4)	3.87 dd (11.2,3.6)	3.73 dd (10.0,4.0)	4.58 dd (10.4,4.0)	3.78 dd (10.4,3.6)	3.83 brd (10.8)	3.74 dd (10.4,3.6)	4.61 dd (10.4,3.6)	3.81 dd (10.4,3.2)
9b	3.74 dd (12.0,6.0)	3.39 dd (10.4,6.4)	4.38 dd (10.4,6.4)	3.55 dd (10.4,7.2)	3.47 dd (11.2,6.0)	3.13 dd (10.0,6.0)	4.00 dd (10.4,6.0)	3.75 dd (10.4,2.0)	3.44 dd (10.8,6.4)	3.12 dd (10.4,6.4)	3.98 dd (10.4,5.2)	3.76 dd (10.4,2.0)
1'	4.48 d (8.0)	4.13 d (7.6)	5.01 d (8.0)	4.23 d (7.6)	4.35 d (7.6)	4.06 (7.6)	4.95 d (7.6)	4.14 d (8.0)	4.32 d (8.0)	4.05 d (7.6)	4.96 d (7.6)	4.15 d (8.0)
2'	3.34 dd (8.0,8.4)	2.96 dd (8.0,7.6) (8.0,8.0)	4.09 dd (8.0,7.6)	3.16 dd (8.0,7.6)	3.32 dd (8.0,7.6)	2.96 dd (8.8,7.6)	4.07 dd (8.0,7.6)	3.16 dd (8.4,8.0)	3.28 dd (8.4,8.0)	2.97 dd (8.4,7.6)	4.08 dd (8.4,7.6)	3.16 dd (8.8,8.0)
3'	3.40 dd (8.4,8.4)	3.13 dd (8.8,8.0) (8.4,8.0)	4.22 dd (8.0,8.0)	3.21 dd (8.0,8.0)	3.38 dd (8.0,8.0)	3.11 dd (8.8,8.8)	4.20 dd (8.8,8.0)	3.22 dd (8.4,8.4)	3.34 dd (8.4,8.4)	3.11 dd (8.4,8.4)	4.21 dd (8.8,8.4)	3.22 dd (8.8,8.4)
4'	3.51 dd (8.8,8.4)	3.05 dd (8.8,8.0) (8.8,8.4)	4.24 dd (8.0,8.0)	3.30 dd (8.0,8.0)	3.49 dd (8.0,8.0)	3.05 dd (8.8,8.0)	4.23 dd (8.8,8.8)	3.29 dd (8.4,8.4)	3.45 dd (8.4,8.4)	3.04 dd (8.4,8.4)	4.25 dd (8.8,8.4)	3.28 dd (8.4,8.4)
5'	3.44 m	3.06 m	3.90 m	3.20 m	3.38 m	3.05 m	3.88 m	3.16 m	3.34 m	3.04 m	3.90 m	3.17 m
6'a	3.92 brd (11.2)	3.64 brd (11.2)	4.47 brd (12.0)	3.78 brd (12.4)	3.88 dd (12.4,1.2)	3.63 brd (11.6)	4.47 dd (12.0,2.0)	3.77 brd (12.0)	3.84 brd (12.0)	3.62 brd (12.0)	4.50 brd (11.6)	3.76 brd (12.0)
6'b	3.72 dd (11.2,6.0)	3.39 dd (11.2,4.8)	4.34 dd (12.0,5.2)	3.60 dd (12.4,5.2)	3.70 dd (12.4,4.4)	3.40 dd (11.6,4.4)	4.32 dd (12.0,5.6)	3.58 dd (12.0,5.2)	3.66 dd (12.4,4.0)	3.40 brd (12.0)	4.35 dd (11.6,5.2)	3.58 dd (12.0,5.2)
OMe	3.90 s	3.73 s	3.65 s	3.81 s	3.90 s	3.73 s	3.70 s	3.80 s	3.85 s	3.72 s	3.77 s	3.79 s

<sup>a</sup>  $^1\text{H}$  NMR data were measured at 400 MHz. Proton coupling constants (*J*) in Hz are given in parentheses.

**Table S15.**  $^{13}\text{C}$  NMR data ( $\delta$ ) for Compounds **14–16** in Different Solvents<sup>a</sup>

no.	<b>14 (Erythro)</b>				<b>15 (Threo)</b>				<b>16 (Threo)</b>			
	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD	D <sub>2</sub> O	DMSO- <i>d</i> <sub>6</sub>	C <sub>5</sub> D <sub>5</sub> N	CD <sub>3</sub> OD
1	135.7	134.1	135.1	135.1	135.6	133.9	135.1	134.5	135.1	133.0	134.2	133.7
2	114.1	111.2	111.9	112.3	113.7	111.1	111.9	111.6	107.4	104.3	105.5	105.2
3	150.2	147.0	148.3	149.3	150.1	146.9	148.3	148.9	150.7	147.4	148.9	149.1
4	147.6	145.3	147.1	147.5	147.3	145.3	147.1	147.2	136.7	134.3	136.7	136.0
5	118.2	114.7	115.8	116.3	118.1	114.7	116.0	115.9	150.7	147.4	148.9	149.1
6	123.2	119.4	120.7	121.4	122.6	119.1	120.4	120.7	107.4	104.3	105.5	105.2
7	<b>76.3</b>	<b>73.6</b>	<b>75.3</b>	<b>76.0</b>	<b>76.6</b>	<b>72.6</b>	<b>74.3</b>	<b>75.4</b>	<b>77.10</b>	<b>72.8</b>	<b>74.7</b>	<b>75.6</b>
8	<b>76.3</b>	<b>73.9</b>	<b>75.6</b>	<b>76.1</b>	<b>76.9</b>	<b>74.0</b>	<b>76.1</b>	<b>76.2</b>	<b>77.17</b>	<b>74.0</b>	<b>76.3</b>	<b>76.2</b>
9	74.0	71.5	73.1	73.0	73.7	70.9	72.3	72.4	73.9	70.8	72.2	72.3
1'	105.7	103.8	105.7	105.5	105.5	103.6	105.5	105.0	105.7	103.6	105.6	105.0
2'	76.1	73.7	75.2	75.7	75.8	73.6	75.2	75.2	76.1	73.6	75.3	75.2
3'	78.5	76.4	78.3	78.4	78.3	76.4	78.3	77.9	78.5	76.5	78.5	77.9
4'	72.5	70.0	71.4	72.0	72.3	69.9	71.5	71.5	72.6	70.0	71.6	71.5
5'	78.7	76.9	78.5	78.5	78.5	76.8	78.4	77.9	78.7	76.8	78.6	77.9
6'	63.6	61.0	62.4	63.1	63.3	61.0	62.5	62.6	63.6	61.0	62.6	62.6
OMe	58.8	55.6	55.7	56.9	58.6	55.6	55.9	56.4	59.3	55.9	56.4	56.8
$\Delta\delta_{\text{C}8-\text{C}7}$	<b>0.0</b>	<b>0.3</b>	<b>0.3</b>	<b>0.1</b>	<b>0.3</b>	<b>1.4</b>	<b>1.8</b>	<b>0.8</b>	<b>0.07</b>	<b>1.2</b>	<b>1.6</b>	<b>0.6</b>

<sup>a</sup>  $^{13}\text{C}$  NMR data were measured at 125 MHz. The assignments were based on DEPT,  $^1\text{H}$ - $^1\text{H}$  COSY, HMQC, and HMBC experiments.